



Identification and expression profile of multiple genes in the anterior kidney of channel catfish induced by modified live *Edwardsiella ictaluri* vaccination

Julia W. Pridgeon^{*}, Craig A. Shoemaker, Phillip H. Klesius

Aquatic Animal Health Research Unit, USDA-ARS, 990 Wire Road, Auburn, AL 36832, USA

ARTICLE INFO

Article history:

Received 10 June 2009

Received in revised form 10 September 2009

Accepted 13 September 2009

Keywords:

Edwardsiella ictaluri

Ictalurus punctatus

Subtractive hybridization

Expression

Modified live vaccination

ABSTRACT

Using PCR-select subtractive cDNA hybridization technique, 57 expressed sequence tags (ESTs) were isolated from 240 clones of a modified live *Edwardsiella ictaluri* vaccinated vs. sham-vaccinated channel catfish anterior kidney subtractive library. The transcription levels of the 57 ESTs in response to *E. ictaluri* vaccination were then evaluated by quantitative PCR (QPCR). Of the 57 ESTs, 43 were induced at least 2-fold higher in all three vaccinated fish compared to unvaccinated control fish. Of the 43 upregulated genes, five were consistently upregulated greater than 10-fold, including two highly upregulated (>20-fold) glycosyltransferase and Toll-like receptor 5. The transcriptional levels of GTPase 1, coatamer protein complex zeta 1, and type II arginine deiminase were consistently induced greater than 10-fold. MHC class I α chain and transposase were upregulated greater than 10-fold in two of the three vaccinated fish. The 43 upregulated genes also included 19 moderately upregulated (3–10-fold) and 17 slightly upregulated (2–3-fold). Our results suggest that subtractive cDNA hybridization and QPCR are powerful cost-effective techniques to identify differentially expressed genes in response to modified live *E. ictaluri* vaccination.

Published by Elsevier B.V.

1. Introduction

Enteric septicemia of catfish (ESC), the most prevalent disease affecting farm-raised channel catfish, *Ictalurus punctatus*, is caused by *Edwardsiella ictaluri*, a facultative intracellular Gram-negative flagellated bacterium akin to phylogenetically related *Salmonella* (Thune et al., 1997; Zhang and Arias, 2007). ESC is generally an acute septicemia that develops very quickly, especially in the temperature range of 22–28 °C. Signs of the disease have been observed within 2 days after immersion challenge and heavy mortalities have been reported as early as 4 days after infection (Newton et al., 1989; Wolters and Johnson, 1994; Thune et al., 1997).

To control ESC, live attenuated *E. ictaluri* vaccines have been developed to protect catfish (Wise et al., 2000; Shoemaker et al., 1999, 2002, 2007; Karsi et al., 2009). Several studies have demonstrated that protective immunity in channel catfish against *E. ictaluri* is largely mediated by cellular immune responses with humoral antibodies having a secondary function (Ellis, 1999; Shoemaker and Klesius, 1997). Like *Salmonella*, *E. ictaluri* can survive and replicate intracellularly (Steele-Mortimer et al., 2000; Skirpstunas and Baldwin, 2002; Thune et al., 2007; Russo et al., 2009), further suggesting that the cellular immune response plays an important role in combating ESC.

The innate immune system is the first line of host defense against pathogens and plays a vital role in maintaining host-microbe homeostasis (Bingle and Craven, 2004). Bony fish have very quick and powerful defense mechanisms to a wide range of pathogens (Bayne et al., 2001; Ellis, 2001). The host immune system recognizes invading pathogens by their

^{*} Corresponding author. Tel.: +1 334 887 3741; fax: +1 334 887 2983.
E-mail address: Julia.Pridgeon@ars.usda.gov (J.W. Pridgeon).

highly conserved pathogen-associated molecular patterns (PAMPs), which are unique to these pathogens and are normally not shared by host cells. Recognition mediated by pattern recognition receptors (PRRs) will then initiate the inflammatory processes (Medzhitov and Janeway, 2002; Magor and Magor, 2001; Hoffman et al., 1999). Many components of PRRs, such as Toll-like receptors (TLRs), are evolutionarily conserved from insects to humans (Kimbrell and Beutler, 2001). TLRs function in innate immunity through recognizing the conserved pathogen-associated molecular patterns (PAMPs) of an invading pathogen and eliciting inflammatory and immune responses (Medzhitov and Janeway, 2000). To date, 13 TLRs (TLR1–TLR13) have been identified in mammals and five TLRs (TLR2, TLR3, TLR5, TLR20, and TLR21) have been reported in channel catfish (Bilodeau and Waldbieser, 2005; Baoprasertkul et al., 2007a,b). The best characterized ligands that TLRs recognize include: (1) lipoproteins by TLR2; (2) dsRNA by TLR3; (3) lipopolysaccharide (LPS) by TLR4; and (4) flagellin by TLR5.

To understand the molecular mechanism involved in host immune response to *E. ictaluri* infection in channel catfish, expression profiles of different genes have been selected and studied. For example, the expression of channel catfish TLR3 and TLR5 in response to *E. ictaluri* challenge through immersion has been studied at 2, 5, 8, and 21 days post-challenge and it has been demonstrated that the expression of TLR3 is significantly upregulated in the head kidney of catfish 2 days after immersion challenge (Bilodeau and Waldbieser, 2005). The expression of channel catfish TLR2 in response to *E. ictaluri* challenge through immersion has been studied at 4, 24, and 72 h post-challenge and it has been demonstrated that TLR2 is downregulated at all time points in the head kidney after immersion challenge (Baoprasertkul et al., 2007a), suggesting that anterior kidney play an important role in the host immune defense system. A recent study has demonstrated that modified live vaccinated catfish are able to control the dispersion of *E. ictaluri* in the anterior kidney (Russo et al., 2009), further suggesting that anterior kidney plays an important role in the immune defense system.

In addition to TLRs, many other genes such as chemokines, antimicrobial peptides, and pro-inflammatory cytokines have been reported to be upregulated in channel catfish challenged by *E. ictaluri* (Peatman et al., 2005, 2006; Chen et al., 2005; Baoprasertkul et al., 2004; Bao et al., 2006; Yeh and Klesius, 2007, 2008). To understand the transcriptional regulation of genes in response to modified live *E. ictaluri* vaccination in the anterior kidney without any preconception of their identities, we used PCR-select suppression subtractive hybridization in this study. Since its first introduction to researchers (Diatchenko et al., 1996), suppression subtractive hybridization has been widely used by researchers in fish innate immunology (Dios et al., 2007; Zhang et al., 2007a,b) and other fields of science (Singh et al., 2008; Zhao et al., 2008; Zhou et al., 2008; Pridgeon et al., 2009) because this technique does not require any previously known genome information for the organism (Sternberg and Gepstein, 2007; Hillmann et al., 2009). We used quantitative PCR (QPCR) to compare the transcriptional levels of genes in vaccinated fish and unvaccinated fish

because QPCR has tremendous sensitivity and requires little or no post-amplification processing (Wong and Medrano, 2005). Furthermore, QPCR is a highly reproducible technique for the gel-free detection and quantification of mRNA (Ashton and Headrick, 2007). Using PCR-select subtractive cDNA hybridization technique, we identified 57 different genes from 240 clones of modified live *E. ictaluri* vaccinated vs. non-vaccinated channel catfish anterior kidney subtractive library. The transcriptional profiles of the 57 genes in response to *E. ictaluri* vaccination and their putative functions in immune defense were discussed in this study.

2. Materials and methods

2.1. Experimental fish, vaccine strain, and vaccination protocol

Channel catfish fry (NWAC-103 strain) were obtained from the USDA-ARS Catfish Genetic Research Unit, Stoneville, MS and maintained at the USDA-ARS-Aquatic Animal Health Unit at Auburn, AL. Fish were maintained in dechlorinated city water in 340 L tanks to ensure that the catfish fingerlings remained naïve to *E. ictaluri* during grow-out. Catfish fingerlings were grown for 8 months before vaccination and were 186 ± 9 g at time of vaccination. Prior to vaccination fish were moved to 208-L flow-through aquaria and acclimated for 14 days. Fish were vaccinated with AQUAVAC-ESC™ following the established protocol from the manufacturer (Intervet/Schering Plough, Millsboro, DE). The vaccine was based on the modified RE-33 *E. ictaluri* developed by Klesius and Shoemaker (1999). Briefly, the vaccine was thawed at 26 °C in a water bath prior to dilution in 18.9 L of water at 26 ± 2 °C. Three channel catfish (*I. punctatus*) were transferred into one half (9.45 L) of the prepared vaccine bath provided with aeration through air stones via an air blower and held for 2 min. After the 2 min exposure time, an equal volume of water was added and the fish were allowed to remain in the vaccine bath for 15 min prior to release into 208-L flow-through tanks with constant aeration at water temperature of 26 ± 2 °C and a 12:12 h light:dark photoperiod. The bacterial concentration was 6.5×10^6 CFU/mL in the vaccine bath. Three additional catfish were sham-vaccinated in immersion water only following the same procedure as above but without the addition of vaccine. All fish were returned to aquarium after vaccination or sham vaccination. At 48 h post-vaccination, fish were euthanized with MS-222 (300 mg/L). The anterior kidney tissue from each fish was collected and flash frozen in liquid nitrogen during collection. All samples were stored at -80 °C until RNA extraction.

2.2. Total RNA extraction and cDNA synthesis

Total RNA was isolated from anterior kidney tissues using TRIzol Reagent (Invitrogen, Carlsbad, CA) following the manufacturer's protocol. All total RNAs were quantified on a Nanodrop ND-1000 spectrophotometer (Nanodrop Technologies, Rockland, DE). Total RNAs were resuspended in distilled water and stored at -80 °C. The first strand cDNAs used for quantitative PCR were synthesized using AMV reverse transcriptase (Invitrogen,

Carlsbad, CA). For subtractive library construction, total RNA were pooled from the three vaccinated or non-vaccinated fish samples. cDNAs were then synthesized from the pooled total RNAs using PCR-selected cDNA Subtraction Kit (Clontech, Palo Alto, CA) as described by the manufacturer. The cDNAs that contain specific transcripts are referred to as “testers” (i.e. from vaccinated fish) and the reference cDNAs are referred to as “drivers” (i.e. from non-vaccinated fish). The double-stranded cDNAs of both testers and drivers were digested with *RsaI* to create smaller blunt-ended fragments to be used as testers or drivers according to the manufacturer's instruction (Clontech, Palo Alto, CA). The tester cDNAs were then subdivided into two portions (A and B) and modified by ligating with cDNA adaptors 1 and 2 (provided by the kit), respectively.

2.3. Construction of subtractive cDNA library

Two-step subtractive hybridizations were performed according to procedures used previously (Pridgeon and Liu, 2003). Briefly, in the first step hybridization, two primary hybridization reactions (A and B) were formed by adding excess amounts of unmodified driver cDNA to separate portions A and B of tester cDNA samples at a 50:1 ratio. The samples were denatured for 2 min at 98 °C and allowed to anneal for 8 h at 68 °C. The remaining single-stranded, adaptor-ligated tester cDNAs were dramatically enriched in each hybridization reaction for overexpressed sequences because non-target cDNAs present in the tester and driver formed hybrids. For the second step hybridization, A and B primary hybridization reaction solutions were mixed together without denaturing. These new hybrids were double-stranded tester molecules with different 5'-ends corresponding to the sequences of two different adaptors. Freshly denatured driver DNA was added to the reaction without denaturing the subtraction mix to further enrich new double-stranded tester molecules that are differentially expressed. After filling in the adapter ends with DNA polymerase, overexpressed sequences (tester cDNA) had different annealing sites on their 3'- and 5'-ends. The molecules were then subjected to suppression subtraction PCR as described by the manufacturer (Clontech, Palo Alto, CA). The PCR products were then cloned into pGEM-T easy vector (Promega, Madison, WI). Plasmids were then transformed into One Shot[®] TOP10 competent cells (Invitrogen, Carlsbad, CA). Transformed cells were then plated out on Luria-Bertani (LB) plates containing ampicillin (100 µg/mL) and X-Gal (5-bromo-4-chloro-3-indolyl-beta-D-galactopyranoside) (40 µg/mL).

2.4. PCR analysis of subtraction efficiency

To evaluate the subtraction efficiency, the relative amount of the constitutively expressed reference gene 18S rRNA was compared in subtracted cDNA and unsubtracted cDNA after 15, 20, 25, 30, 35, and 40 cycles of PCR. Primers used for the amplification of the 18S rRNA gene was 18S-F (5'-ATGGCCGTCTTCTAGTTGGTG-3') and 18S-R (5'-TAGG-TAGCACACGCTGATCG-3'). The two primers were designed based on channel catfish 18S small subunit ribosomal RNA gene sequence (GenBank accession no. BE469353).

2.5. Plasmid DNA isolation and sequencing

From the library, a total of 240 white colonies were present and subsequently picked to grow overnight in LB broth in the presence of ampicillin (100 µg/mL) in the Innova[™] 4000 Incubator Shaker (New Brunswick Scientific, Edison, NJ) at 37 °C and 235 rpm settings, respectively. Plasmid DNAs were isolated with QIAprep Spin Miniprep Kit (QIAGEN, Valencia, CA). Plasmid DNAs were then digested by *EcoRI* at 37 °C for 1 h and subjected to 1% agarose gel electrophoresis. Plasmid DNAs that contained inserts were then sent to USDA-ARS Mid South Genomic Laboratories in Stoneville, MS for sequencing with an ABI 3730 Genetic Analyzer (Applied Biosystems, Foster City, CA). Sequences were then analyzed using the National Center for Biotechnology Information (NCBI) BLAST program to search for sequence homologies.

2.6. Primer design and quantitative PCR

Sequencing results of different clones were used to design gene-specific primers by using Primer3 program (http://frodo.wi.mit.edu/cgi-bin/primer3/primer3_www.cgi). QPCR was performed on an Applied Biosystems 7300 Fast Real-Time PCR System (Applied Biosystems, Foster City, CA). For each cDNA sample, channel catfish 18S ribosomal RNA primers were included as an internal control to normalize the variation of cDNA amount. All QPCR was performed on an Applied Biosystems 7000 Real-Time PCR System (ABI, Foster City, CA) using Platinum[®] SYBR[®] Green qPCR SuperMix-UDG with ROX (Invitrogen, Carlsbad, CA) in a total volume of 12.5 µL. The QPCR mixture consisted of 1 µL of cDNA, 0.5 µL of 5 µM gene-specific forward primer, 0.5 µL of 5 µM gene-specific reverse primer and 10.5 µL of 1× SYBR Green SuperMix. The QPCR thermal cycling parameters were 50 °C for 2 min, 95 °C for 10 min followed by 40 cycles of 95 °C for 15 s and 60 °C for 1 min. All QPCR was run in duplicate for each cDNA sample and three fish cDNA samples were analyzed by QPCR.

2.7. QPCR data analysis

The relative transcriptional levels of different genes were determined by subtracting the cycle threshold (C_t) of the sample by that of the 18S rRNA, the calibrator or internal control, as per the formula: $\Delta C_t = C_t(\text{sample}) - C_t(\text{calibrator})$. The relative expression level of the specific gene in *E. ictaluri* RE-33 vaccinated fish compared to that in non-vaccinated fish was then calculated by the formula $2^{-\Delta\Delta C_t}$ where $\Delta\Delta C_t = \Delta C_t(\text{vaccinated}) - \Delta C_t(\text{non-vaccinated})$ as described previously (Pridgeon et al., 2009). Data were analyzed by analysis of variance (ANOVA) using SigmaStat statistical analysis software (Systat Software, San Jose, CA).

3. Results

3.1. Evaluation of subtraction efficiency

The subtraction efficiency was evaluated by PCR analysis of the constitutively expressed gene 18S rRNA.




Fig. 1. Evaluation of subtraction efficiency. PCR analysis of the reduction of 18S rRNA abundance was performed using subtracted (lanes 1–6) and unsubtracted (lanes 7–12) cDNA samples as templates for 15 cycles (lanes 1 and 7), 20 cycles (lanes 2 and 8), 25 cycles (lanes 3 and 9), 30 cycles (lanes 4 and 10), 35 cycles (lanes 5 and 11), and 40 cycles (lanes 6 and 12). m: 100 bp DNA ladder.

As shown in Fig. 1, the amount of the 18S rRNA was significantly decreased after subtraction. Obvious bands were seen after 30 cycles in unsubtracted cDNAs but only after 35 and 40 cycles in subtracted cDNAs. The abundance of 18S rRNA was theoretically reduced by 2^5 – 2^{10} times, suggesting that cDNAs specific for vaccinated fish were enriched for about 2^5 – 2^{10} times by suppression hybridization.

3.2. Characteristics of the subtractive cDNA library

A total of 240 clones were obtained from the anterior kidney subtractive library. Of the 240 clones, 182 contained inserts. Sequencing results revealed that these 182 ESTs represented 57 different genes (Table 1). All ESTs listed in Table 1 have been deposited in the GenBank dbEST under accession numbers GO898766–GO898822. The most frequently detected clones were glycosyltransferase ($n=8$), neutrophil cytosolic factor 2 ($n=4$), Rab11A ($n=4$), caldendrin ($n=4$), and LAMP5 ($n=4$). Out of the 57 ESTs identified from the subtractive library, 19 shared homology with deposited *Danio rerio* proteins, seven shared homology with deposited *Ictalurus punctatus* proteins, and 6 shared homology with deposited *Salmo salar* proteins (Table 1). The biggest insert size was 842 bp (GO898807) and the smallest insert size was 119 bp (GO898781). The average insert size of the 57 ESTs was 387 bp (Table 1).

3.3. Expression profiling of the 57 ESTs in catfish after *E. ictaluri* vaccination

To determine whether the expression levels of the 57 ESTs isolated from the subtractive library were upregulated in *E. ictaluri* vaccinated catfish, gene-specific primers for the 57 ESTs were designed (Table 2) for relative QPCR experiments. QPCR results revealed that 43 ESTs were induced at least 2-fold higher in all three vaccinated fish compared to that in unvaccinated control fish (Table 3). Of the 43 upregulated genes, five were consistently upregulated greater than 10-fold (Table 3), including two highly upregulated (>20-fold) genes: glycosyltransferase (GTFase) and Toll-like receptor 5 (TLR5) (Fig. 2A). The transcriptional levels of GTPase 1, coatomer protein complex zeta 1 (CPCZ1), and type II arginine deiminase (ADI2) were consistently induced greater than 10-fold (Table 3). MHC class I α chain and transposase were

Table 1
List of genes isolated from the *E. ictaluri* vaccinated vs. non-vaccinated catfish anterior kidney subtractive cDNA library.

| No. | Accession no. | Protein homology | Protein accession no. | Organism | Identities (%) | Score bits | e value | Insert size (bp) |
|-----|---------------|--|-----------------------|---------------------------------|----------------|------------|---------|------------------|
| 1 | GO898766 | Glycosyltransferase | CAL56729 | <i>Ostreococcus taur</i> | 56% | 33.9 | 4.9 | 291 |
| 5 | GO898767 | DNA topoisomerase I | ZP_02330037 | <i>Paenibacillus</i> | 33% | 33.5 | 6.2 | 432 |
| 6 | GO898768 | Ring finger 144B | NP_001107142 | <i>Xenopus tropicalis</i> | 88% | 134 | 2e–30 | 317 |
| 7 | GO898769 | NADH dehydrogenase 1 alpha subcomplex 4 | XP_001234601 | <i>Gallus gallus</i> | 89% | 86 | 1e–13 | 338 |
| 20 | GO898770 | Propionyl Coenzyme A carboxylase β | CAQ13492 | <i>Danio rerio</i> | 89% | 163 | 5e–39 | 243 |
| 26 | GO898771 | Neutrophil cytosolic factor 2 | BAF73667 | <i>Cyprinus carpio</i> | 76% | 126 | 5e–28 | 248 |
| 27 | GO898772 | Surface antigen BspA-like | XP_001315236 | <i>Trichomonas vaginalis</i> | 27% | 33.1 | 8.1 | 475 |
| 28 | GO898773 | Ras-related protein Rab 11A | NP_001134031 | <i>Salmo salar</i> | 95% | 144 | 3e–33 | 235 |
| 31 | GO898774 | Kruppel-like protein 1 | XP_001944423 | <i>Acyrtosiphon pisum</i> | 36% | 33.1 | 8.2 | 208 |
| 32 | GO898775 | c-mos | AAL55336 | <i>Bachia flavescens</i> | 48% | 33.1 | 8.1 | 290 |
| 33 | GO898776 | Metacaspase-like protein | XP_680261 | <i>Plasmodium berghei</i> | 41% | 33.9 | 6.0 | 525 |
| 34 | GO898777 | Coatomer protein complex, zeta 1 | NP_571583 | <i>Danio rerio</i> | 100% | 204 | 2e–51 | 316 |
| 41 | GO898778 | Slc3a2 protein | AAI59203 | <i>Danio rerio</i> | 61% | 102 | 1e–20 | 249 |
| 42 | GO898779 | No homology | – | – | – | – | – | 296 |
| 58 | GO898780 | Cytochrome b | AAI79027 | <i>Ictalurus punctatus</i> | 97% | 253 | 5e–66 | 417 |
| 70 | GO898781 | Complement C4a | BAB03284 | <i>Cyprinus carpio</i> | 51% | 41.2 | 0.031 | 119 |
| 77 | GO898782 | Splicing factor 3a, subunit 3 | CAK04939 | <i>Danio rerio</i> | 99% | 224 | 2e–57 | 427 |
| 78 | GO898783 | Glucose phosphate isomerase a | AAH83507 | <i>Danio rerio</i> | 91% | 367 | 7e–100 | 739 |
| 81 | GO898784 | Adenosine deaminase-related growth factor | XP_643866 | <i>Dictyostelium discoideum</i> | 51% | 36.2 | 0.97 | 471 |
| 83 | GO898785 | Huntingtin interacting protein K | XP_001922634* | <i>Danio rerio</i> | 95% | 182 | 1e–44 | 553 |
| 85 | GO898786 | Tumor necrosis factor, alpha-induced protein 2 | XP_697839 | <i>Danio rerio</i> | 57% | 156 | 1e–36 | 688 |

Table 1 (Continued)

| No. | Accession no. | Protein homology | Protein accession no. | Organism | Identities (%) | Score bits | e value | Insert size (bp) |
|-----|---------------|--|-----------------------|---------------------------------|----------------|------------|---------|------------------|
| 86 | G0898787 | Lysosomal-associated transmembrane protein 5 | XP_001923185 | <i>Danio rerio</i> | 56% | 55.1 | 2e–06 | 221 |
| 92 | G0898788 | Alpha-tubulin isotype M-alpha-2 | NP_001098596 | <i>Danio rerio</i> | 99% | 235 | 1e–60 | 353 |
| 93 | G0898789 | NADH dehydrogenase subunit 3 | YP_913699 | <i>Hypentelium nigricans</i> | 88% | 89.4 | 1e–16 | 372 |
| 101 | G0898790 | Caldendrin | XP_001511181 | <i>Ornithorhynchus anatinus</i> | 89% | 92.8 | 2e–17 | 669 |
| 107 | G0898791 | Beta thymosin-like | NP_001124169 | <i>Danio rerio</i> | 92% | 78.6 | 2e–13 | 272 |
| 120 | G0898792 | Granulin 1 | AAI53544 | <i>Danio rerio</i> | 63% | 97.1 | 5e–19 | 300 |
| 121 | G0898793 | CD83 antigen precursor | ACI67585 | <i>Salmo salar</i> | 29% | 44.7 | 0.008 | 760 |
| 122 | G0898794 | Leukocyte cell-derived chemotaxin 2 | NP_001041520 | <i>Danio rerio</i> | 74% | 114 | 2e–24 | 212 |
| 125 | G0898795 | TrkA domain-containing protein | YP_001528048 | <i>Desulfococcus oleovorans</i> | 30% | 35.4 | 2.3 | 556 |
| 126 | G0898796 | Ribosomal protein L5a | AAK95128 | <i>Ictalurus punctatus</i> | 100% | 244 | 1e–63 | 363 |
| 127 | G0898797 | Neurobeachin-like 1 | XP_001070093 | <i>Rattus norvegicus</i> | 78% | 89.0 | 1e–16 | 169 |
| 128 | G0898798 | Multisubunit cleavage/polyadenylation specificity factor subunit A | XP_730560 | <i>Plasmodium yoelii</i> | 26% | 34.3 | 7.6 | 646 |
| 134 | G0898799 | lysozyme g | ACN09851 | <i>Salmo salar</i> | 73% | 214 | 2e–54 | 510 |
| 136 | G0898800 | CD45 precursor | AAV65851 | <i>Ictalurus punctatus</i> | 95% | 349 | 4e–95 | 488 |
| 137 | G0898801 | Very large inducible GTPase 1 | XP_6840863 | <i>Danio rerio</i> | 43% | 105 | 2e–21 | 352 |
| 139 | G0898802 | SET translocation B | NP_958876 | <i>Danio rerio</i> | 98% | 341 | 3e–92 | 690 |
| 148 | G0898803 | Tubulin alpha 8 like | CAQ14258 | <i>Danio rerio</i> | 97% | 213 | 6e–54 | 317 |
| 151 | G0898804 | Eukaryotic translation initiation factor 3, subunit E | NP_001135167 | <i>Salmo salar</i> | 98% | 164 | 2e–39 | 241 |
| 159 | G0898805 | Rapunzel 2 | NP_001138713 | <i>Danio rerio</i> | 68% | 129 | 8e–29 | 405 |
| 165 | G0898806 | Pre-mRNA processing factor 8 | XP_002197327 | <i>Taeniopygia guttata</i> | 98% | 199 | 7e–50 | 280 |
| 169 | G0898807 | Prostate stem cell antigen precursor-like/UPAR | ABD85498 | <i>Ictalurus punctatus</i> | 84% | 247 | 1e–63 | 842 |
| 174 | G0898808 | DNA or RNA helicase of superfamily II | ZP_01235811 | <i>Vibrio angustum S14</i> | 65% | 33.1 | 8.3 | 373 |
| 177 | G0898809 | Tubulin, alpha, ubiquitous | ABY89801 | <i>Callithrix jacchus</i> | 99% | 237 | 3e–61 | 353 |
| 178 | G0898810 | Predicted protein | XP_002288130 | <i>Thalassiosira pseudonana</i> | 35% | 47.0 | 6e–04 | 376 |
| 179 | G0898811 | Protein-arginine deiminase type II-like | ABE98234 | <i>Oreochromis mossambicus</i> | 77% | 133 | 6e–30 | 286 |
| 184 | G0898812 | Uroporphyrinogen decarboxylase | AAI08076 | <i>Danio rerio</i> | 84% | 233 | 3e–58 | 318 |
| 195 | G0898813 | Tropomyosin alpha-3 chain | NP_001136186 | <i>Salmo salar</i> | 92% | 113 | 8e–24 | 532 |
| 205 | G0898814 | MHC class I alpha chain | AAD08649 | <i>Ictalurus punctatus</i> | 87% | 120 | 4e–26 | 188 |
| 210 | G0898815 | Matrix metalloproteinase-9 | ABO86718 | <i>Ictalurus punctatus</i> | 95% | 85.9 | 1e–15 | 126 |
| 212 | G0898816 | Sulfotransferase family, cytosolic, 2B, member 1a | XP_541518 | <i>Canis familiaris</i> | 28% | 34.3 | 3.8 | 370 |
| 217 | G0898817 | Beta-actin | ABY62772 | <i>Squalius alburnoides</i> | 100% | 304 | 1e–81 | 452 |
| 218 | G0898818 | Toll-like receptor 5 | ABF74618 | <i>Ictalurus punctatus</i> | 96% | 186 | 8e–46 | 434 |
| 219 | G0898819 | Solute carrier family 25, member 3 isoform 3 | XP_002189856 | <i>Taeniopygia guttata</i> | 100% | 42.4 | 0.014 | 455 |
| 223 | G0898820 | Transposase | ABV31710 | <i>Salmo salar</i> | 90% | 224 | 2e–57 | 366 |
| 224 | G0898821 | Glutamic pyruvate transaminase 2 | NP_001092227 | <i>Danio rerio</i> | 94% | 113 | 5e–24 | 165 |
| 239 | G0898822 | NAD-dependent deacetylase sirtuin-5 | ACI33678 | <i>Salmo salar</i> | 68% | 34.7 | 2.8 | 243 |

Table 2
Gene-specific primers used in QPCR.

| Clone no. | Accession no. | Forward primer (5'–3') | Reverse primer (5'–3') |
|-----------|---------------|-------------------------|------------------------|
| 1 | G0898766 | GCTTCGAGGTCAGATGAGA | AGTGTTCATGGCCTCTGC |
| 5 | G0898767 | CAGGGTCGACTGTAAAGCA | GTGCGGGAACGACTGTTTAT |
| 6 | G0898768 | TTCTTCTGTGCCAGGGTTT | GGGGACATTTTCCTGAGACA |
| 7 | G0898769 | CGGCCAGACTTCTAAATCCA | TGCGGTTAGCTGTTTTCT |
| 20 | G0898770 | TCACCATCATCACCAGGAAG | TGATTCTCCTTTCCCTGAA |
| 26 | G0898771 | GGGTTTAACCGGAGCATCTT | ATTGAGCTGAGCTGCTGTT |
| 27 | G0898772 | TTGTGGAATGCCTCTGACTG | TTTTGCCGTAGACCGAAGAC |
| 28 | G0898773 | GCCATGACGGGTAGAGAAGA | CAGATCTGGGCTTTGATCGT |
| 31 | G0898774 | ATGCCTTTTCTACCACCTC | AACTCGTGCAAGGCTGAAAT |
| 32 | G0898775 | GTTTCATGCAGCATTCACACC | CCACAACACATCAGGTTCCA |
| 33 | G0898776 | TGGAATGCAACGAATCA | AAAGAAGGCAAGCAGAATGTG |
| 34 | G0898777 | AGAAAAGGCCCTCCATGTTT | TTGCTCGAAGCCTTACTGT |
| 41 | G0898778 | ATCCTTTGCTTGGCATTGAG | TTTTTCAGCCAGTAGGCACA |
| 42 | G0898779 | TTAATGAGTCGGTGCTGCTG | AATCCTGCCCTCCTTCAGTT |
| 58 | G0898780 | GGAGGTTGGGAGAGAAAAGG | TCCGAGCAACACTACTCCAC |
| 70 | G0898781 | CGGGAAGTGCTACAGCG | CCTAAAGCGCGTCTCAG |
| 77 | G0898782 | ACGTCACCCAGATCGAAGAC | AAGGGGTTCTCAAGGTGTCT |
| 78 | G0898783 | CTCCAGAGCATCACCCTCA | CTGCTCAATCTCAGCATCCA |
| 81 | G0898784 | TGCAGATGCTGCTTAAATG | AGCTGTGGCTTTTATTGGA |
| 83 | G0898785 | TCGAGTTGCGACTTGGAAAC | TTGTCAGTGAATCAAAAGC |
| 85 | G0898786 | TCATGTATGACCAGCTCA | CTTGATGGGGTGATAGACA |
| 86 | G0898787 | TCGCGTGCTTGTGTTATG | CAGGTGTCCATTGCTGTTA |
| 92 | G0898788 | TCAGATCGTGCTCTCCATCA | CTCAAAGCAGGCATTTGTGA |
| 93 | G0898789 | AGAAGGGTGTAAGCGGGAGT | AAAAGCTCTCCCCCTACGAA |
| 101 | G0898790 | CCAGCCAGCAGAAAGTAAGG | AGGGAAGCCATGAGGAAACT |
| 107 | G0898791 | GACAAACCAATCTCG | CCTGCTTCTCCTGTTCG |
| 120 | G0898792 | ACATGCTGCAGGTCTCCTTT | TCTGAATGGTGTGCTCTGC |
| 121 | G0898793 | AAACGACAAGAGACCACAGAA | CAGCCCTTTGACTCTGGAAT |
| 122 | G0898794 | AAACAACGCCATCAACAATG | GTTGCTGCTTGTAGCTTTTC |
| 125 | G0898795 | GTGACAGCAGGGGTTTTGT | CGGCTTTAGGGTGGATATGA |
| 126 | G0898796 | AGTGCCTTCACGTGCTACCT | GTTCAGGCCCAGAATGTGTT |
| 127 | G0898797 | CACCAAGGAGAGCACTACA | TCTCCTGCAGACCTGAGA |
| 128 | G0898798 | AAACACGTGCAGGAGGAACT | CGTGCTTCGTTAGCTTTTC |
| 134 | G0898799 | GAGCCTGGAACAGTGAGGAG | GTGGTGCTCTGTCCATTTT |
| 136 | G0898800 | GTTCCGCAGTTGCAGATTTT | TGACGTTACTTTCCAGGTG |
| 137 | G0898801 | TCCATGAGCACAGTGAGAG | AAGGCTCATCTTGGGGTTTT |
| 139 | G0898802 | CTCTGCTTGGTGAGGAGGAC | AGTCTTTCCAGCCTCCAT |
| 148 | G0898803 | TGATGGCAATATGCCAAGTG | CTGCATCTTCTCCAGAG |
| 151 | G0898804 | CTGAGCGCTGTCAAAGTCAA | ACCTAACACAGCGGTCATC |
| 159 | G0898805 | CACCTTCTCCTGACCTGCTC | AGTCTTTTGGGAGGCCACT |
| 165 | G0898806 | TCCTCCAGATAACCCACAGG | ATGATGGTCTTCTGCCATC |
| 169 | G0898807 | TGCATCAGTGCATCAGTTCA | GGACGATCATGAGGAGGAAA |
| 174 | G0898808 | AAACGCTACGGAGCTTTGAA | CACCTGGTTCATAGAGGACA |
| 177 | G0898809 | TCAGATCGTGCTCTCCATCA | CTCAAAGCAGGCATTTGTGA |
| 178 | G0898810 | TCAGATCGTGCTCTCCATCA | CTCAAAGCAGGCATTTGTGA |
| 179 | G0898811 | TGAGACGTGCTCTTGTCTG | TCTCACAGTCAAGGTTCCA |
| 184 | G0898812 | CAAGGGTCAATGTGTTCAATC | GACATCCGACCGAGAGTAAA |
| 195 | G0898813 | CGCCTTATCATATGCGGTCT | CCCCACCTTCTACCCACTTT |
| 205 | G0898814 | CACCAGGAATAAAATTTCCAGA | CTGTGCTCGGTGTTCCAGT |
| 210 | G0898815 | ACATCTTATTAGTAAGAGGGTCC | ACTACTGGAAGTCTCAAATCG |
| 212 | G0898816 | CAACATGAGAGCGAAGTGGA | AGCAGATGGACAGCACCTCT |
| 217 | G0898817 | CTCGAAGTCAAGGGCAACAT | CGTGATGACTCTGGTGATG |
| 218 | G0898818 | TTGGAAGCGGTACAAATCCT | ACCCGAGGTTGAATAATCC |
| 219 | G0898819 | CAGAAACAGCTTTGCACCTG | CTCATTCCTCTGTCATCAC |
| 223 | G0898820 | ACCGTCAAACTCTGGGTGAG | GCAACTCGAACCTTCAGCTC |
| 224 | G0898821 | AAACAGCACTGCCAGC | ATACACCTCATCGGCCA |
| 239 | G0898822 | CCAAAAGGAGGCACACTTG | GCCAAAGATGCCAGAATACC |

upregulated greater than 10-fold in two of the three vaccinated fish. The transcription level of NAD-dependent deacetylase sirtuin-5 was induced greater than 8-fold in two of the three vaccinated fish. The transcriptional levels of c-mos, adenosine deaminase-related growth factor, huntingtin interacting protein K, and tumor necrosis factor α -induced protein 2 were consistently upregulated greater than 5-fold (Fig. 2B). Other genes that were consistently

induced greater than 3-fold and included the following: Kruppel-like protein 1, matrix metalloproteinase-9, DNA topoisomerase I, propionyl coenzyme A carboxylase β , netrophil cytosolic factor 2, surface antigen BspA-like, complement C4a, granulin 1, glutamic pyruvate transaminase 2, rapunzel 2, prostate stem cell antigen precursor-like/plasminogen activator urokinase receptor, DNA or RNA helicase of superfamily II, a predicted protein with

Table 3Modified live *E. ictaluri* vaccine induced expression of the 57 genes.

| Gene/clone | Accession no. | Putative protein | Vaccine induced expression ($2^{\Delta\Delta C_T}$) (fold) | | |
|------------|---------------|--|--|--------------------------|--------------------------|
| | | | $2^{\Delta\Delta C_T-1}$ | $2^{\Delta\Delta C_T-2}$ | $2^{\Delta\Delta C_T-3}$ |
| 1 | G0898766 | DNA topoisomerase I | 72.05 | 101.48 | 32.11 |
| 5 | G0898767 | Ring finger 144B | 5.35 | 5.68 | 3.45 |
| 6 | G0898768 | NADH dehydrogenase 1 alpha subcomplex 4 | 2.46 | 2.27 | 2.98 |
| 7 | G0898769 | Propionyl coenzyme A carboxylase β | 1.79 | 2.07 | 3.16 |
| 20 | G0898770 | Neutrophil cytosolic factor 2 | 3.84 | 3.31 | 3.61 |
| 26 | G0898771 | Surface antigen BspA-like | 3.53 | 5.15 | 4.77 |
| 27 | G0898772 | Ras-related protein Rab 11A | 3.07 | 3.38 | 4.79 |
| 28 | G0898773 | Kruppel-like protein 1 | 1.85 | 2.63 | 5.22 |
| 31 | G0898774 | c-mos | 4.25 | 4.6 | 9.06 |
| 32 | G0898775 | Metacaspase-like protein | 6.71 | 6.45 | 7.39 |
| 33 | G0898776 | Coatamer protein complex, zeta 1 | 2.94 | 2.27 | 3.24 |
| 34 | G0898777 | Slc3a2 protein | 10.93 | 18.25 | 18.25 |
| 41 | G0898778 | No homology | 2.67 | 3.49 | 5.56 |
| 42 | G0898779 | Cytochrome <i>b</i> | 2.53 | 2.91 | 3.96 |
| 58 | G0898780 | Complement C4a | 3.14 | 3.04 | 2.92 |
| 70 | G0898781 | Splicing factor 3a, subunit 3 | 3.79 | 4.94 | 6.77 |
| 77 | G0898782 | Glucose phosphate isomerase a | 2.12 | 2.6 | 2.34 |
| 78 | G0898783 | Adenosine deaminase-related growth factor | 1.72 | 2.14 | 3.07 |
| 81 | G0898784 | Huntingtin interacting protein K | 5.35 | 6.13 | 6.8 |
| 83 | G0898785 | Tumor necrosis factor, alpha-induced protein 2 | 5.03 | 5.58 | 8.11 |
| 85 | G0898786 | Lysosomal-associated transmembrane protein 5 | 7.84 | 6.06 | 11.16 |
| 86 | G0898787 | Alpha-tubulin isotype M-alpha-2 | 2.89 | 3.73 | 4.91 |
| 92 | G0898788 | NADH dehydrogenase subunit 3 | 1.74 | 2.11 | 3.75 |
| 93 | G0898789 | Caldendrin | 5.28 | 4.13 | 2.82 |
| 101 | G0898790 | Beta thymosin-like | 1.83 | 3.14 | 3.31 |
| 107 | G0898791 | Granulin 1 | 0.91 | 1.17 | 1.56 |
| 120 | G0898792 | CD83 antigen precursor | 3.21 | 4.36 | 5.12 |
| 121 | G0898793 | Leukocyte cell-derived chemotaxin 2 | 2.61 | 3.47 | 8.25 |
| 122 | G0898794 | TrkA domain-containing protein | 3.68 | 3.12 | 6.08 |
| 125 | G0898795 | Ribosomal protein L5a | 1.42 | 2.2 | 2.48 |
| 126 | G0898796 | Neurobeachin-like 1 | 1.02 | 1.09 | 1.46 |
| 127 | G0898797 | Multisubunit cleavage/polyadenylation specificity factor subunit A | 2.91 | 4.61 | 4.55 |
| 128 | G0898798 | Lysozyme g | 2.79 | 4.87 | 6.45 |
| 134 | G0898799 | CD45 precursor | 2.53 | 2.17 | 3.17 |
| 136 | G0898800 | Very large inducible GTPase 1 | 1.55 | 9.03 | 3.59 |
| 137 | G0898801 | SET translocation B | 15.81 | 30.8 | 25.11 |
| 139 | G0898802 | Tubulin alpha 8 like | 1.51 | 4.08 | 3.4 |
| 148 | G0898803 | Eukaryotic translation initiation factor 3, subunit E | 2.77 | 8.85 | 8.97 |
| 151 | G0898804 | Rapunzel 2 | 2.22 | 5.84 | 3.58 |
| 159 | G0898805 | Pre-mRNA processing factor 8 | 3.87 | 9.32 | 5.92 |
| 165 | G0898806 | Prostate stem cell antigen precursor-like/UPAR | 2.5 | 5.17 | 4.86 |
| 169 | G0898807 | DNA or RNA helicase of superfamily II | 3.32 | 6.41 | 4.89 |
| 174 | G0898808 | Tubulin, alpha, ubiquitous | 3.66 | 6.04 | 6.34 |
| 177 | G0898809 | Predicted protein | 2.18 | 3.22 | 4.52 |
| 178 | G0898810 | Protein-arginine deiminase type II-like | 3.12 | 3.39 | 3.95 |
| 179 | G0898811 | Uroporphyrinogen decarboxylase | 11.15 | 17.88 | 11.43 |
| 184 | G0898812 | Tropomyosin alpha-3 chain | 0.54 | 0.46 | 1.38 |
| 195 | G0898813 | MHC class I alpha chain | 3.16 | 6.52 | 2.7 |
| 205 | G0898814 | Matrix metalloproteinase-9 | 17.53 | 26.54 | 4.71 |
| 210 | G0898815 | Sulfotransferase family, cytosolic, 2B, member 1a | 4.2 | 4.71 | 4.21 |
| 212 | G0898816 | Beta-actin | 1.14 | 1.11 | 1.58 |
| 217 | G0898817 | Toll-like receptor 5 | 1.12 | 1.16 | 1.52 |
| 218 | G0898818 | Solute carrier family 25, member 3 isoform 3 | 31.61 | 43.26 | 21.63 |
| 219 | G0898819 | Transposase | 1.35 | 2.24 | 2.23 |
| 223 | G0898820 | Glutamic pyruvate transaminase 2 | 20.25 | 7.73 | 10.48 |
| 224 | G0898821 | NAD-dependent deacetylase sirtuin-5 | 5 | 6.77 | 3.59 |
| 239 | G0898822 | | 19.46 | 8.52 | 3.64 |

unknown function, and leucocyte derived chemotaxin 2 (LECT2). Seventeen genes slightly upregulated 2–3-fold included the following: ringer finger 144B, metacaspase-like protein, slc3a2 protein, cytochrome *b*, splicing factor 3a3, lysosomal-associated transmembrane protein 5 (LAMP5), NADH dehydrogenase subunit 3, CD83 antigen

precursor, neurobeachin-like 1 multisubunit cleavage/polyadenylation specificity factor subunit A, lysozyme g, tubulin alpha 8, eukaryotic translation initiation factor 3E, pre-mRNA processing factor 8, tubulin alpha ubiquitous, tropomyosin alpha-3 chain, and one novel protein (Fig. 2C). Nine genes induced greater than 2-fold in two of the three




Fig. 2. Transcriptional regulation of the 57 genes isolated from the channel catfish head kidney subtractive library. (A) Transcription levels of 5 highly upregulated (>10-fold) genes induced by *E. ictaluri* vaccination; (B) transcription levels of 21 moderately upregulated (3 < x < 10-fold) genes induced by *E. ictaluri* vaccination; (C) transcription levels of 17 slightly upregulated (2 < x < 3-fold) genes induced by *E. ictaluri* vaccination; (D) transcription levels of 14 genes not significantly induced by *E. ictaluri* vaccination (<2-fold). The relative transcriptional levels of different genes were determined by subtracting the cycle threshold (C_t) of the sample by that of the 18S rRNA, the calibrator or internal control, as per the formula: $\Delta C_t = C_t (\text{sample}) - C_t (\text{calibrator})$. The relative expression level of the specific gene in *E. ictaluri* RE-33 vaccinated fish compared to that in non-vaccinated fish was then calculated by the formula $2^{-\Delta\Delta C_t}$ where $\Delta\Delta C_t = \Delta C_t (\text{vaccinated}) - \Delta C_t (\text{non-vaccinated})$. Data are presented as means \pm SEM. Differences were considered statistically significant when $p < 0.05$ and represented by an asterisk.

vaccinated fish included the following: Ras-related protein Rab-11A, NADH dehydrogenase 1 alpha subcomplex 4, glucose phosphate isomerase a, alpha-tubulin isotype M-alpha-2, caldendrin, TrkA domain-containing protein, CD45 precursor, SET translocation (myeloid leukemia-associated) B, and solute carrier family 25, member 3 isoform 3. Five genes that were induced less than 2-fold in all three vaccinated fish were the following: ribosomal protein L5a, sulfotransferase family cytosolic 2B member 1a, beta-actin, beta thymosin-like, and uroporphyrinogen decarboxylase (Fig. 2D).

3.4. Classification of the *E. ictaluri* vaccination-upregulated genes

The 57 genes isolated from the subtractive library were classified in terms of their putative functions (Table 4). Half of the genes identified were either involved in immune-response or metabolism (Fig. 3). The major portion (28%) of the 57 genes identified were immune-related genes, including MHC class I alpha chain, surface antigen BspA-like protein/NK-lysin type 3, complement C4a, leukocyte cell-derived chemotaxin 2 (LECT2), CD83 antigen pre-

Table 4Putative function of *E. ictaluri* vaccination-upregulated genes.

| Category | Protein | Putative function related to infection |
|--|---|---|
| Immune-related protein (16) | Glycosyltransferase | Immune response-T cell apoptosis |
| | Similar to very large inducible GTPase 1 | Mediate in innate and adaptive immunity |
| | MHC class I alpha chain | Antiviral immunity |
| | Surface antigen BspA-like/NK-lysin type 3 | Cell attachment and invasion |
| | Complement C4-1 | Acute infection |
| | Granulin 1 | Phagocytosis |
| | Prostate stem cell antigen precursor-like/urokinase plasminogen activator receptor (uPAR or PLAUR) | Modulate the development of protective immunity |
| | Leukocyte-derived chemotaxin 2 (LECT2) | Infection induced gene |
| | CD83 antigen precursor | Dendritic cell maturation |
| | Lysozyme g | Bactericidal activity |
| | c-mos | Phagocytosis |
| | Matrix metalloproteinase-9 | Host defense |
| | Neutrophil cytosolic factor 2 | Phagocytosis |
| | Toll-like receptor 5 | Innate immunity |
| | CD45 precursor | Type I interferon production |
| | Similar to tumor necrosis factor, alpha-induced protein 2 | Innate immunity |
| Metabolism-related protein (12) | NAD-dependent deacetylase sirtuin-5 | Cellular metabolism |
| | Similar to sulfotransferase family, cytosolic, 2B, member 1 isoform a | Mitochondria burst/Metabolism |
| | NADH dehydrogenase (ubiquinone) 1 alpha subcomplex, 4, | Phagocytosis/oxidative stress response |
| | Cytochrome b | Mitochondria burst/Bacteria pathogenesis |
| | Splicing factor 3a, subunit 3 | Xenobiotics detoxification |
| | Propionyl Coenzyme A carboxylase, beta polypeptide | Metabolism |
| | Glutamic pyruvate transaminase (alanine aminotransferase) 2 | metabolism |
| | Solute carrier family 25 (mitochondrial carrier; phosphate carrier), member 3 isoform 3 | metabolism |
| | Uroporphyrinogen decarboxylase | metabolism |
| | NADH dehydrogenase subunit 3 | metabolism |
| | Glucose phosphate isomerase a | metabolism |
| | Protein-arginine deiminase type II-like | metabolism |
| Apoptosis-related protein (6) | Slc3a2 protein | Apoptosis-related/CD98HC; Secretory pathways |
| | Beta thymosin-like | Apoptosis |
| | similar to huntingtin interacting protein K | Apoptosis |
| | Metacaspase-like protein | Apoptosis |
| | Similar to Kruppel-like protein 1 | Apoptosis |
| | Ring finger 144B | Apoptosis |
| Cell structure, growth and maintenance (8) | Beta-actin | Cell structure |
| | Similar to alpha-tubulin isotype M-alpha-2 | Cell structure |
| | Tubulin, alpha 8 like | Cell structure |
| | Tubulin, alpha, ubiquitous (predicted) | Cell structure |
| | Tropomyosin alpha-3 chain | Actin-related |
| | SET translocation (myeloid leukemia-associated) B | Nucleosome assembly |
| | ribosomal protein L5a | |
| | adenosine deaminase-related growth factor | Cell proliferation |
| Endocytosis, vesicular, and lysosome trafficking (3) | Similar to lysosomal-associated transmembrane protein 5 (lysosomal-associated multitransmembrane protein 5) | Lysosome trafficking |
| | (Retinoic acid-inducible E3 protein) | |
| | coatamer protein complex, zeta 1 | Vesicle trafficking |
| | Ras-related protein Rab-11A | Endocytosis and trafficking |
| Transcription and translation related protein (3) | PRP8 pre-mRNA processing factor 8 homolog | Transcription regulation |
| | Eukaryotic translation initiation factor 3, subunit E, a | Translation |
| | Multisubunit cleavage/polyadenylation specificity factor subunit A | Transcription |
| Bacterial proteins (3) | DNA topoisomerase I | bacteria |
| | DNA or RNA helicase of superfamily II | bacteria |
| | TrkA domain-containing protein | Bacteria |

Table 4 (Continued)

| Category | Protein | Putative function related to infection |
|-----------------------------------|-------------------------|--|
| Signal transduction (1) | Caldendrin | Calcium signaling |
| Functionally unknown proteins (5) | No homology | unknown |
| | Unnamed protein product | unknown |
| | Transposase | unknown |
| | Rapunzel 2 | unknown |
| | Predicted protein | unknown |

cursor, lysozyme g, CD45 precursor, and Toll-like receptor 5, glucose phosphate isomerase a, and TNF alpha-induced protein 2 (TNFAIP2). Another major group (21%) of the identified genes was related to metabolism, NADH dehydrogenase subunit 3 and cytochrome *b*. Six genes (10%) were related to apoptosis, including huntingtin interacting protein K and ring finger 144B. Eight genes (14%) were related to cell growth and maintenance, including beta-actin and alpha-tubulin. Three genes (5%) were related to endocytosis and lysosome trafficking, including lysosomal-associated transmembrane protein 5 (LAMP5) and Ras-related protein Rab-11A. Three genes (5%) were related to gene regulation, including pre-mRNA processing factor 8 and eukaryotic translation initiation factor 3. One gene (2%), caldendrin, was involved in calcium signaling. The functions of five genes (9%) are currently unknown.

4. Discussion

Enteric septicemia of catfish (ESC) is the most prevalent disease affecting farm-raised channel catfish (Wagner et al., 2002) and live attenuated *E. ictaluri* vaccines have been shown to be efficacious in protecting catfish through immersion (Wise et al., 2000; Shoemaker et al., 1999, 2002, 2007). However, it is not clear whether vaccination will induce consistent molecular responses in channel catfish after exposure to *E. ictaluri* vaccine. Therefore, the purpose of this study was to isolate upregulated genes in channel catfish after modified live *E. ictaluri* vaccination without any preconception of their identities. We sampled fish at 48 h post-vaccination because kinetic studies have

revealed that pathogen loads are peaked on day 2 in channel catfish and that Toll-like receptor 3 is significantly induced on day 2 in channel catfish after immersion challenge with *E. ictaluri* (Bilodeau & Waldbieser, 2005). Using subtractive cDNA hybridization and QPCR, Toll-like receptor 5 (TLR5) was identified to be consistently upregulated greater than 20-fold in all three vaccinated catfish. Toll-like receptor 5 is a well-characterized receptor in both mammals and fish that recognizes flagellin of bacteria (Gewirtz et al., 2001; Tsujita et al., 2004). The expression of channel catfish TLR5 in response to virulent strain of *E. ictaluri* has been demonstrated to be significantly upregulated in the anterior kidney and in the liver following ESC infection (Bilodeau and Waldbieser, 2005; Bilodeau et al., 2006; Peatman et al., 2008). The identification of TLR5 from the subtractive library without any preconception of TLR5's identity confirms that suppression subtractive hybridization is a powerful technique to identify upregulated genes in one mRNA population vs. another.

EST G0898766 was consistently upregulated greater than 20-fold in all three vaccinated fish. At protein level, G0898766 shared 56% identity with glycosyltransferase of *Ostreococcus taur*, a free-living eukaryote. However, the *e* value of 4.9 in the blast search suggested that G0898766 might be a functionally unknown protein. The function of glycosyltransferase is to catalyze the transfer of sugar moieties from activated donor molecules to specific acceptor molecules, forming glycosidic bonds. The acceptor molecule can be a lipid, a protein, a heterocyclic compound, or another carbohydrate residue. Recently, a putative glucosyltransferase designated biofilm-associated




Fig. 3. Classification of the differentially expressed genes identified from the subtractive library. Pie charts representing the distribution of the 57 identified genes according to their putative biological function.

glycolipid synthesis A (*bgsA*) in bacterium *Enterococcus faecalis* has been demonstrated to synthesize DGlcdAG, a glycolipid and lipoteichoic acid precursor involved in biofilm accumulation, adherence to host cells, and virulence *in vivo* (Theilacker et al., 2009), suggesting that this EST might be a virulence gene in *E. ictaluri*. It is also possible that GO898766 is a host immune-related gene since it has been suggested that the expression of glycosyltransferase can modulate cell death during T cell development and function through controlling susceptibility to galectin-1 (Galvan et al., 2000). The function of galectin-1 is to induce apoptosis of immature thymocytes and activate T cells (Galvan et al., 2000). 5'- and 3'-rapid amplification of cDNA ends will be merited to understand the true identity of ESTGO898766.

EST GO898801 was upregulated greater than 15-fold in all three vaccinated fish. GO898801 shared 43% identity with a very large inducible GTPase 1 of zebra fish *Danio rerio* at protein level (e value = $2e-21$). Very large inducible GTPase 1 has been reported to be inducible by interferons, important players in both innate immunity and adaptive immunity (Klamp et al., 2003). Recently, it has been demonstrated that the expression of guanylate-binding protein-1 (GBP-1), an interferon inducible large GTPase, is upregulated in colonic epithelia of individuals with inflammatory bowel disease (Schnoor et al., 2009), suggesting that this very large inducible GTPase 1 might be a specific response to *E. ictaluri* vaccination and/or infection.

EST GO898777 was upregulated greater than 10-fold in all three vaccinated fish. GO898777 shared 100% identity with coatamer protein complex zeta 1 of zebra fish *Danio rerio* at protein level (e value = $2e-51$). Coatamer protein complex is involved in endocytosis and vesicle trafficking (Maier et al., 2001). It has been demonstrated that coatamer protein complex is necessary for the maintenance and/or localization of a critical pool of elements of the phagocytic machinery (Hackam et al., 2001), suggesting that the upregulation of coatamer protein complex zeta 1 might play an important role in phagocytosis of *E. ictaluri* in the channel catfish to survive ESC.

EST GO898811 was upregulated greater than 11-fold in all three vaccinated fish. GO898811 shared 77% identity with type II arginine deiminase of Mozambique tilapia *Oreochromis mossambicus* at protein level (e value = $6e-30$). The function of arginine deiminase (ADI) is to catalyze the irreversible hydrolysis of arginine to citrulline and ammonia (Shirai et al., 2001). ADI has been shown to be able to inhibit lipopolysaccharide (LPS)-induced upregulation of inducible nitric oxide synthase and the production of nitric oxide in murine macrophages in dose-dependent manner (Kim et al., 2007), suggesting that ADI upregulation in the channel catfish might be a specific response to *E. ictaluri* infection and/or vaccination.

EST GO898814 was upregulated greater than 17-fold in two of the three vaccinated fish. GO898814 shared 87% identity with major histocompatibility complex (MHC) class I alpha chain of channel catfish *Ictalurus punctatus* at protein level (e value = $4e-26$). MHC plays an important role in the immune system. Proteins encoded by the MHC are expressed on the surface of cells, displaying both self

antigens (peptide fragments from the cell itself) and nonself antigens (e.g. fragments of invading microorganisms) to T cell that has the capacity to kill or co-ordinate the killing of pathogens and infected or malfunctioning cells. Upregulation of MHC class I alpha chain has been reported in the liver of blue catfish 3 days post-challenge by *E. ictaluri* (Peatman et al., 2008), suggesting that the upregulation of MHC class I alpha chain might play an important role in the host immune response to *E. ictaluri* infection and/or vaccination.

EST GO898822 was upregulated greater than 8-fold in two of the three vaccinated fish. GO898822 shared 68% identity with NAD-dependant deacetylase sirtuin-5 protein of salmon *Salmo salar*. However, the e-value was only 2.8, suggesting that sirtuin-5 sequence in channel catfish might be very different from that in salmon. The enzymes of the sirtuin family of nicotinamide-adenine-dinucleotide-dependent protein deacetylases are emerging key players in nuclear and cytosolic signaling, but also in mitochondrial regulation and aging. Sirtuin-5 has been demonstrated to be able to deacetylates carbamoyl phosphate synthetase 1 and regulates the urea cycle in mice (Nakagawa et al., 2009). Sirtuin 5 has also been demonstrated to be able to deacetylate cytochrome c, a protein of the mitochondrial intermembrane space with a central function in oxidative metabolism as well as apoptosis initiation (Schlicker et al., 2008). The upregulation of sirtuin 5 in channel catfish in response to *E. ictaluri* vaccination suggests that sirtuin 5 might also play an important role in host immune defense.

EST GO898772 was upregulated greater than 3-fold in all three vaccinated fish. GO898772 shared 27% identity with surface antigen BspA-like protein of eukaryotic pathogen *Trichomona vaginalis* at protein level (e value = 8.1). It has been reported that epithelial cell attachment and invasion by bacteria *Tannerella forsythia* are dependent on the BspA protein (Inagaki et al., 2006). However, at nucleotide level, GO898772 shared 76% identity with NK-lysin type 3 gene of the channel catfish (e value = $5e-38$). NK-lysin is a 9-kDa polypeptide that was originally isolated from porcine intestinal tissue based on its antibacterial activity. NK lysine has been reported to be able to bind LPS from *Escherichia coli*, *Pseudomonas aeruginosa*, and different strains of *Salmonella enterica* and lyse lymphoma cells (Andersson et al., 1999), suggesting that NK-lysin type 3 may be a natural LPS-binding protein that may participate in the endogenous defense response in the channel catfish in response to *E. ictaluri* vaccination.

EST GO898781 was upregulated greater than 3-fold in all three vaccinated fish. GO898781 shared 51% identity with complement C4a of common carp *Cyprinus carpio* (e value = 0.031). The complement system is an important humoral defense mechanism that plays a vital role against microbial agents, inflammatory response control, and immunocomplex clearance. Classical complement pathway activation is antibody-dependent. The C4 component participates in the initial step of activation, and C4 expression is determined by 2 pairs of allotypes: C4a and C4b (Samano et al., 2004; Rambach et al., 2008). It has been reported that C4a levels was significantly higher in

acute Lyme disease patients than in tick bite and healthy control groups (Shoemaker et al., 2008). In *E. ictaluri* challenged blue catfish liver, C4b expression has been reported to be upregulated for 3.9-fold compared to non-challenged control (Peatman et al., 2008), suggesting that the upregulation of C4a in channel catfish might serve as a molecular marker for *E. ictaluri* infection and/or vaccination.

EST G0898792 was upregulated greater than 3-fold in all three vaccinated fish. G0898792 shared 63% identity with granulins-1 protein of zebra fish *Danio rerio* (e value = $5e-19$). The granulins are small proteins of about 6 kDa that are derived from a larger precursor of approximately 590 amino acids (Bhandari et al., 1992, Plowman et al., 1992). Both the 6-kDa peptide and the intact precursor have been found to modulate cell growth (Bateman and Bennett, 1998). It has been reported that equine granulins E is able to kill *Streptococcus zooepidemicus* and sustain a greater than 99.8% decrease in CFU per milliliter after a 2-h exposure to 100 $\mu\text{g/mL}$ (approximately 15 μM) of granulins E (Couto et al., 1992), suggesting that the upregulation of granulins-1 in vaccinated channel catfish might be involved in host immune defense against *E. ictaluri* vaccination and/or infection.

EST G0898807 was upregulated 3–6-fold in all three vaccinated fish. G0898807 shared 84% identities with prostate stem cell antigen precursor-like protein of channel catfish (e value = $1e-63$). G0898807 also shared 39% identity with urokinase plasminogen activator receptor of zebra fish with an e value of $2e-08$. The urokinase plasminogen activator receptor (uPAR) is expressed at the cell surface of inflammatory cells and plays an important role in neutrophil migration. It has been reported that uPAR is crucially involved in host defense through phagocytosis during *E. coli* induced acute pyelonephritis in mice (Roelofs et al., 2006), suggesting that upregulation of uPAR might play an important role in host defense against *E. ictaluri* vaccination and/or infection.

EST G0898794 was upregulated 3–6-fold in all three vaccinated fish. G0898794 shared 74% identity with leukocyte cell-derived chemotaxin 2 (LECT2) of zebra fish (e value = $2e-24$). LECT2 was initially isolated as a possible chemotactic factor for neutrophils (Yamagoe et al., 1996) and subsequently demonstrated to be involved in liver regeneration (Saito et al., 2004), carcinogenesis (Ovejero et al., 2004) and NK cell homeostasis (Saito et al., 2004). Recently, it has been reported that LECT2 is induced 51-fold by *Staphylococcus aureus* and 1344-fold by *Aeromonas salmonicida* infection in zebra fish (Lin et al., 2007), suggesting that upregulation of LECT2 might play an important role in anti-infection and anti-inflammation.

EST G0898793 was upregulated 2.6–8-fold in three vaccinated fish. G0898793 shared 29% identity with CD83 antigen precursor of atlantic salmon *Salmo salar* (e value = 0.008). CD83, an evolutionarily well-conserved highly glycosylated type 1 transmembrane glycoprotein, is a highly specific marker for activated dendritic cells (DCs) in human. The function of CD83 is to regulate lymphocyte maturation, activation and homeostasis (Bre-

loer and Fleischer, 2008). It has been reported that CD83 is rapidly upregulated on murine B cells, reaching maximal expression 6 h after either Toll-like receptor (TLR) engagement by lipopolysaccharide (LPS) or B cell receptor (BCR) ligation (Kretschmer et al., 2007). It has been reported that monocyte-derived dendritic cells is able to recognize and respond to the essential viral glycoproteins, leading to the upregulation of CD40, CD83, and CD86 in response to herpes simplex virus-1 infection (Reske et al., 2008), suggesting that upregulation of CD83 might play important functional role in the channel catfish in response to *E. ictaluri* infection and/or vaccination.

EST G0898799 was upregulated 2–3-fold in three vaccinated fish. BLAST homology search revealed that G0898799 shared 73% identity with lysozyme g of atlantic salmon *Salmo salar* (e value = $2e-54$). Lysozyme functions as a crucial biodefense effector against the infection of bacterial pathogens in innate immunity. It has been reported that the expression of g-type lysozyme is significantly upregulated in the vaccinated Atlantic cod, *Gadus morhua* after intraperitoneal vaccination of heat-killed *Listonella anguillarum* (Caipang et al., 2008), suggesting that lysozyme g upregulation might be a specific immune response to *E. ictaluri* vaccination.

Other genes that were upregulated greater than 3-fold in all three vaccinated fish include ESTs that shared identities with c-mos, adenosine deaminase-related growth factor, kruppel-like protein, matrix metalloproteinase-9, and neutrophil cytosolic factor 2. The proto-oncogene c-mos regulates macrophage differentiation in a dose-dependent manner (Kurata et al., 1989). Adenosine deaminase-related growth factor plays an important role in cell proliferation (Zurovec et al., 2002). High steady state expression of kruppel-like protein has been reported in Ag-specific CD8⁺ memory T cells, critical for controlling viral as well intracellular and parasitic infections (Grayson et al., 2001). Matrix metalloproteinase-9 (MMP-9) is involved in the migration of inflammatory cells across the extracellular matrix, as well as tissue remodeling (Delclaux et al., 1996; Opdenakker et al., 2001). MMP-9 was first identified in neutrophils, but can also be expressed by various other cell types such as monocytes/macrophages, lymphocytes, and endothelial cells. MMP-9 is not produced constitutively, but needs a trigger to be expressed (Opdenakker et al., 2001). However, Yeh and Klesius (2008) have suggested that MMP-9 is constitutively expressed in restricted tissues including the head kidney of channel catfish. It has been demonstrated that LPS, the major constituent of the outer cell wall of Gram-negative bacteria and the principal mediator of inflammatory responses to these pathogens, is able to induce the release of MMP-9 by neutrophils and monocytes in vitro (Masure et al., 1991; Opdenakker et al., 1991). Moreover, in mice, *E. coli* LPS administration has led to a quick release of MMP-9 into the circulation, with peak values as soon as 1 h after injection (Dubois et al., 2002), suggesting that the upregulation of MMP-9 in the channel catfish might be specifically induced by *E. ictaluri* vaccination. Neutrophil cytosolic factor 2 (NCF2) is a protein encodes p67^{phox}, a subunit of the multiprotein enzyme complex NADPH oxidase. NCF2 has been shown to be responsible for the

generation of superoxides. The phagocyte NADPH oxidase has a crucial role in innate immunity by specifically reducing molecular oxygen to superoxide. Superoxide anions give rise to numerous toxic reactive oxygen species that are used as microbicidal agents against pathogens, contributing to the respiratory burst typically seen in phagocytic cells. Upon activation by opsonized microbes or inflammatory mediators, p67^{phox} will be translocated to the membrane to form an active enzyme complex with flavocytochrome *b*₅₅₈ to regulate electron transfer from NADPH to flavin adenine dinucleotide. Once activated, superoxide is released into the phagocytic vacuole or into the extracellular space (Mizuki et al., 1998; Han et al., 1998; Nisimoto et al., 1999). The upregulation of these five genes induced by *E. ictaluri* vaccination suggests that they might play important roles in the immune defense system in channel catfish in response to *E. ictaluri* vaccination and/or infection.

In summary, 57 different genes were isolated from *E. ictaluri* vaccinated vs. non-vaccinated channel catfish anterior kidney. Of the 57 ESTs, 43 were induced at least 2-fold higher in all three vaccinated fish compared to that in unvaccinated control fish. Of the 43 upregulated genes, five were consistently upregulated greater than 10-fold, including two highly upregulated (>20-fold) glycosyltransferase and Toll-like receptor 5. The transcriptional levels of GTPase 1, coatamer protein complex zeta 1, and type II arginine deiminase were consistently induced greater than 10-fold. MHC class I α chain and transposase were upregulated greater than 10-fold in two of the three vaccinated fish. The 43 upregulated genes also included 19 moderately upregulated (3–10-fold) and 17 slightly upregulated (2–3-fold). Our results suggest that subtractive cDNA hybridization and QPCR are powerful cost-effective techniques to identify differentially expressed genes in response to bacteria challenge. Significant upregulation of multiple genes induced by modified live *E. ictaluri* vaccination suggests that they might play important roles in the host defense by priming the immune system for response to *E. ictaluri* vaccination and/or infection.

Acknowledgments

We thank Drs. Benjamin LaFrentz (USDA-ARS), Zhanjiang Liu and Huseyin Kucuktas (Auburn University) for critical reviews of the manuscript. We thank Dr. Brian Scheffler and Fanny Liu (USDA-ARS-Catfish Genetics Research Unit) for their excellent sequencing work. We thank Stacey A. LaFrentz and Paige Mumma (USDA-ARS) for their excellent technical support. We also thank the management team of the Aquatic Animal Health Research Unit for daily care and management of the fish. This study was supported by the USDA/ARS CRIS project #6420-32000-020-00D. The use of trade, firm, or corporate names in this publication is for the information and convenience of the reader. Such use does not constitute an official endorsement or approval by the United States Department of Agriculture or the Agricultural Research Service of any product or service to the exclusion of others that may be suitable.

References

- Andersson, M., Girard, R., Cazenave, P., 1999. Interaction of NK lysis, a peptide produced by cytolytic lymphocytes, with endotoxin. *Infect. Immun.* 67, 201–205.
- Ashton, K.J., Headrick, J.P., 2007. Quantitative (real-time) RT-PCR in cardiovascular research. *Methods Mol. Biol.* 366, 121–143.
- Bao, B., Peatman, E., Xu, P., Baoprasertkul, B., Wang, G., Liu, Z., 2006. Characterization of 23 CC chemokine genes and analysis of their expression in channel catfish (*Ictalurus punctatus*). *Dev. Comp. Immunol.* 30, 783e96.
- Baoprasertkul, P., Peatman, E., Abernathy, J., Liu, Z., 2007a. Structural characterization and expression analysis of toll-like receptor 2 gene from catfish. *Fish Shellfish Immunol.* 22, 418–426.
- Baoprasertkul, P., Peatman, E., Chen, L., He, C., Kucuktas, H., Li, P., Wang, S., Somridhivej, B., Dunham, R., Liu, Z., 2004. Sequence analysis and expression of a CXC chemokine in resistant and susceptible catfish after infection of *Edwardsiella ictaluri*. *Dev. Comp. Immunol.* 28, 769–780.
- Baoprasertkul, P., Xu, P., Peatman, E., Kucuktas, H., Liu, Z., 2007b. Divergent Toll-like receptors in catfish (*Ictalurus punctatus*): TLR5S, TLR20, TLR21. *Fish Shellfish Immunol.* 23, 1218–1230.
- Bateman, A., Bennett, H.P., 1998. Granulins: the structure and function of an emerging family of growth factors. *J. Endocrinol.* 158, 145–151.
- Bayne, C.J., Gerwick, L., Fujiki, K., Nakao, M., Yano, T., 2001. Immune relevant (including acute phase) genes identified in the livers of rainbow trout, *Oncorhynchus mykiss*, by means of suppression subtractive hybridization. *Dev. Comp. Immunol.* 25, 205e17.
- Bhandari, V., Palfree, R.G.E., Bateman, A., 1992. Isolation and sequence of the granulins precursor cDNA from human bone marrow reveals tandem cysteine-rich granulins domains. *Proc. Natl. Acad. Sci. U.S.A.* 89, 1715–1719.
- Bilodeau, A.L., Peterson, B.C., Bosworth, B.G., 2006. Response of toll-like receptors, lysozyme, and IGF-I in back-cross hybrid F1 male blue channel X female channel catfish challenged with virulent *Edwardsiella ictaluri*. *Fish Shellfish Immunol.* 20, 29–39.
- Bilodeau, A.L., Waldbieser, G.C., 2005. Activation of TLR3 and TLR5 in channel catfish exposed to virulent *Edwardsiella ictaluri*. *Dev. Comp. Immunol.* 29, 713–721.
- Bingle, C.D., Craven, C.J., 2004. Meet the relatives: a family of BPI- and LBP-related proteins. *Trends Immunol.* 25, 53–55.
- Breloer, M., Fleischer, B., 2008. CD83 regulates lymphocyte maturation, activation and homeostasis. *Trends Immunol.* 29, 186–194.
- Caipang, C.M., Hynes, N., Puangkaew, J., Brinchmann, M.F., Kiron, V., 2008. Intraperitoneal vaccination of Atlantic cod, *Gadus morhua* with heat-killed *Listonella anguillarum* enhances serum antibacterial activity and expression of immune response genes. *Fish Shellfish Immunol.* 24, 314–322.
- Chen, L., He, C., Baoprasertkul, P., Xu, P., Li, P., Serapion, J., Waldbieser, G., Wolters, W., Liu, Z., 2005. Analysis of a catfish gene resembling interleukin-8: cDNA cloning, gene structure, and expression after infection with *Edwardsiella ictaluri*. *Dev. Comp. Immunol.* 29, 135–142.
- Couto, M.A., Harwig, S.S.L., Cullor, J.S., Hughes, J.P., Lehrer, R.I., 1992. Identification of eNAP-1, an antimicrobial peptide from equine neutrophils. *Infect. Immun.* 60, 3065–3071.
- Delclaux, C., Delacourt, C., D'Ortho, M.P., Boyer, V., Lafuma, C., Harf, A., 1996. Role of gelatinase B and elastase in human polymorphonuclear neutrophil migration across basement membrane. *Am. J. Respir. Cell Mol. Biol.* 14, 288–295.
- Diatchenko, L., Lau, Y.F., Campbell, A.P., Chenchik, A., Moqadam, F., Huang, B., Lukyanov, S., Lukyanov, K., Gurskaya, N., Sverdlov, E.D., Siebert, P.D., 1996. Suppression subtractive hybridization: a method for generating differentially regulated or tissue-specific cDNA probes and libraries. *Proc. Natl. Acad. Sci. U.S.A.* 93, 6025–6030.
- Dios, S., Poisa-Beiro, L., Figueras, A., Novoa, B., 2007. Suppression subtraction hybridization (SSH) and macroarray techniques reveal differential gene expression profiles in brain of sea bream infected with nodavirus. *Mol. Immunol.* 44, 2195–2204.
- Dubois, B., Starckx, S., Pagenstecher, A., Oord, J., Arnold, B., Opdenakker, G., 2002. Gelatinase B deficiency protects against endotoxin shock. *Eur. J. Immunol.* 32, 2163–2171.
- Ellis, A.E., 1999. Immunity to bacteria in fish. *Fish Shellfish Immunol.* 9, 291–308.
- Ellis, A.E., 2001. Innate host defense mechanisms of fish against viruses and bacteria. *Dev. Comp. Immunol.* 25, 827e39.
- Galvan, M., Tsuboi, S., Fukuda, M., Baum, L.G., 2000. Expression of a specific glycosyltransferase enzyme regulates T cell death mediated by galectin-1. *J. Biol. Chem.* 275, 16730–16737.

- Gewirtz, A.T., Navas, T.A., Lyons, S., Godowski, P.J., Madara, J.L., 2001. Cutting edge: bacterial flagellin activates basolaterally expressed TLR5 to induce epithelial proinflammatory gene expression. *J. Immunol.* 167, 1882–1885.
- Grayson, J.M., Murali-Krishna, K., Altman, J.D., Ahmed, R., 2001. Gene expression in antigen-specific CD8⁺ T cells during viral infection. *J. Immunol.* 166, 795–799.
- Hackam, D.J., Botelho, R.J., Sjolín, C., Rotstein, O.D., Robinson, J.M., Schreiber, A.D., Grinstein, S., 2001. Indirect role for COPI in the completion of Fcγ receptor-mediated phagocytosis. *J. Biol. Chem.* 276, 18200–18208.
- Han, C.H., Freeman, J.L., Lee, T., Motalebi, S.A., Lambeth, J.D., 1998. Regulation of the neutrophil respiratory burst oxidase: identification of an activation domain in p67^{phox}. *J. Biol. Chem.* 273, 16663–16668.
- Hillmann, A., Dunne, E., Kenny, D., 2009. cDNA amplification by SMART-PCR and suppression subtractive hybridization (SSH)-PCR. *Methods Mol. Biol.* 496, 223–243.
- Hoffman, J.A., Kafatos, F.C., Janeway, C.A., Ezekowitz, R.A.B., 1999. Phylogenetic perspectives in innate immunity. *Science* 284, 1313–1318.
- Inagaki, S., Onishi, S., Kuramitsu, H.K., Sharma, A., 2006. Porphyromonas gingivalis vesicles enhance attachment, and the leucine-rich repeat BspA protein is required for invasion of epithelial cells by “Tannerella forsythia”. *Infect. Immun.* 74, 5023–5028.
- Karsi, A., Gölşoy, N., Corb, E., Dumpala, P.R., Lawrence, M.L., 2009. High-throughput bioluminescence-based mutant screening strategy for identification of bacterial virulence genes. *Appl. Environ. Microbiol.* 75, 2166–2175.
- Kim, J.E., Hur, H.J., Lee, K.W., Lee, H.J., 2007. Anti-inflammatory effects of recombinant arginine deiminase originating from *Lactococcus lactis* ssp. *lactis* ATCC 7962. *J. Microbiol. Biotechnol.* 17, 1491–1497.
- Kimbrell, D.A., Beutler, B., 2001. The evolution and genetics of innate immunity. *Nat. Rev. Genet.* 2, 256–267.
- Klump, T., Boehm, U., Schenk, D., Pfeffer, K., Howard, J.C., 2003. A giant GTPase, very large inducible GTPase-1, is inducible by IFNs. *J. Immunol.* 171, 1255–1265.
- Klesius, P.H., Shoemaker, C.A., 1999. Development and use of modified live *Edwardsiella ictaluri* vaccine against enteric septicemia of catfish. *Adv. Vet. Med.* 41, 523–537.
- Kretschmer, B., Lüthje, K., Guse, A.H., Ehrlich, S., Koch-Nolte, F., Haag, F., Fleischer, B., Breloer, M., 2007. CD83 modulates B Cell function in vitro: increased IL-10 and reduced Ig secretion by CD83Tg B cells. *PLoS ONE* 2, e755.
- Kurata, N., Akiyama, H., Taniyama, T., Marunouchi, T., 1989. Dose-dependent regulation of macrophage differentiation by mos mRNA in a human monocytic cell line. *EMBO J.* 8, 457–463.
- Lin, B., Chen, S., Cao, Z., Lin, Y., Mo, D., Zhang, H., Gu, J., Dong, M., Liu, Z., Xu, A., 2007. Acute phase response in zebrafish upon *Aeromonas salmonicida* and *Staphylococcus aureus* infection: striking similarities and obvious differences with mammals. *Mol. Immunol.* 44, 295–301.
- Magor, B.G., Magor, K.E., 2001. Evolution of effectors and receptors of innate immunity. *Dev. Comp. Immunol.* 25, 651–682.
- Maier, A.G., Webb, B., Ding, M., Bremser, M., Carrington, M., Clayton, C., 2001. The coatome of *Trypanosoma brucei*. *Mol. Biochem. Parasitol.* 115, 55–61.
- Masure, S., Proost, P., Van Damme, J., Opdenakker, G., 1991. Purification and identification of 91-kDa neutrophil gelatinase: release by the activating peptide interleukin-8. *Eur. J. Biochem.* 198, 391–398.
- Medzhitov, R., Janeway, C.A., 2000. The toll receptor family and microbial recognition. *Trends Microbiol.* 8, 452–456.
- Medzhitov, R., Janeway Jr., C.A., 2002. Decoding the patterns of self and nonself by the innate immune system. *Science* 296, 298–300.
- Mizuki, K., Kadomatsu, K., Hata, K., Ito, T., Fan, Q.W., Kage, Y., Fukumaki, Y., Sakaki, Y., Takeshige, K., Sumimoto, H., 1998. Functional modules and expression of mouse p40^{phox} and p67^{phox}, SH3-domain-containing proteins involved in the phagocyte NADPH oxidase complex. *Eur. J. Biochem.* 251, 573–582.
- Nakagawa, T., Lomb, D.J., Haigis, M.C., Guarente, L., 2009. SIRT5 Deacetylates carbamoyl phosphate synthetase 1 and regulates the urea cycle. *Cell* 137, 560–570.
- Newton, J.C., Wolfe, L.G., Grizzle, J.M., Plumb, J.A., 1989. Pathology of experimental enteric septicemia in channel catfish, *Ictalurus punctatus* (Rafinesque), following immersion-exposure to *Edwardsiella ictaluri*. *J. Fish Dis.* 12, 335–347.
- Nisimoto, Y., Motalebi, S., Han, C.H., Lambeth, J.D., 1999. The p67^{phox} activation domain regulates electron flow from NADPH to flavin in flavocytochrome b558. *J. Biol. Chem.* 274, 22999–23005.
- Opdenakker, G., Masure, S., Grillet, B., Van Damme, J., 1991. Cytokine-mediated regulation of human leukocyte gelatinases and role in arthritis. *Lymphokine Cytokine Res.* 10, 317–324.
- Opdenakker, G., Van den Steen, P.E., Van Damme, J., 2001. Gelatinase B: a tuner and amplifier of immune functions. *Trends Immunol.* 22, 571–579.
- Ovejero, C., Cavard, C., Périanin, A., Hakvoort, T., Vermeulen, J., Godard, C., Fabre, M., Chafey, P., Suzuki, K., Romagnolo, B., Yamagoe, S., Perret, C., 2004. Identification of the leukocyte cell-derived chemotaxin 2 as a direct target gene of beta-catenin in the liver. *Hepatology* 40, 167–176.
- Peatman, E., Bao, B., Baoprasertkul, P., Liu, Z., 2005. In silico identification and expression analysis of 12 novel CC chemokines in catfish. *Immunogenetics* 57, 409–419.
- Peatman, E., Bao, B., Peng, X., Baoprasertkul, P., Brady, Y., Liu, Z., 2006. Catfish CC chemokines: genomic clustering, duplications, and expression after bacterial infection with *Edwardsiella ictaluri*. *Mol. Genet. Genomics* 275, 297–309.
- Peatman, E., Terhune, J., Baoprasertkul, P., Xu, P., Nandi, S., Wang, S., Somridhivej, B., Kucuktas, H., Li, P., Dunham, R., Liu, Z., 2008. Microarray analysis of gene expression in the blue catfish liver reveals early activation of the MHC class I pathway after infection with *Edwardsiella ictaluri*. *Mol. Immunol.* 45, 553–566.
- Plowman, G.D., Green, J.M., Neubauer, M.G., Buckley, S.D., McDonald, V., Todaro, G.J., Shoyab, M., 1992. The epithelin precursor encodes two proteins with opposing activities on epithelial cell growth. *J. Biol. Chem.* 267, 13073–13078.
- Pridgeon, J.W., Becnel, J.J., Clark, G.G., Linthicum, K.J., 2009. Permethrin induces overexpression of multiple genes in *Aedes aegypti*. *J. Med. Entomol.* 46, 580–587.
- Pridgeon, J.W., Liu, N., 2003. Overexpression of the cytochrome c oxidase subunit I gene associated with a pyrethroid resistant strain of German cockroaches, *Blattella germanica* (L.). *Insect Biochem. Mol. Biol.* 33, 1043–1048.
- Rambach, G., Würzner, R., Speth, C., 2008. Complement: an efficient sword of innate immunity. *Contrib. Microbiol.* 15, 78–100.
- Reske, A., Pollara, G., Krummenacher, C., Katz, D.R., Chain, B.M., 2008. Glycoprotein-dependent and TLR2-independent innate immune recognition of herpes simplex virus-1 by dendritic cells. *J. Immunol.* 180, 7525–7536.
- Roelofs, J.J., Rouschop, K.M., Teske, G.J., Claessen, N., Weening, J.J., van der Poll, T., Florquin, S., 2006. The urokinase plasminogen activator receptor is crucially involved in host defense during acute pyelonephritis. *Kidney Int.* 70, 1942–1947.
- Russo, R., Shoemaker, C.A., Panangala, V.S., Klesius, P.H., 2009. *In vitro* and *in vivo* interaction of macrophages from vaccinated and non-vaccinated channel catfish (*Ictalurus punctatus*) to *Edwardsiella ictaluri*. *Fish Shellfish Immunol.* 26, 543–552.
- Samano, E.S., Ribeiro, L., de M., Gorescu, R.G., Rocha, K.C., Grumach, A.S., 2004. Involvement of C4 allotypes in the pathogenesis of human diseases. *Rev. Hosp. Clin. Fac. Med. Sao Paulo* 59, 138–144.
- Saito, T., Okumura, A., Watanabe, H., Asano, M., Ishida-Okawara, A., Sakagami, J., Sudo, K., Hatano-Yokoe, Y., Bezradica, J.S., Joyce, S., Abo, T., Iwakura, Y., Suzuki, K., Yamagoe, S., 2004. Increase in hepatic NKT cells in leukocyte cell-derived chemotaxin 2-deficient mice contributes to severe concanavalin A-induced hepatitis. *J. Immunol.* 173, 579–585.
- Schlicker, C., Gertz, M., Papatheodorou, P., Kachholz, B., Becker, C.F., Steegborn, C., 2008. Substrates and regulation mechanisms for the human mitochondrial sirtuins Sirt3 and Sirt5. *J. Mol. Biol.* 382, 790–801.
- Schnoor, M., Betanzos, A., Weber, D.A., Parkos, C.A., 2009. Guanylate-binding protein-1 is expressed at tight junctions of intestinal epithelial cells in response to interferon-gamma and regulates barrier function through effects on apoptosis. *Mucosal Immunol.* 2, 33–42.
- Shirai, H., Blundell, T.L., Mizuguchi, K., 2001. A novel superfamily of enzymes that catalyze the modification of guanidino groups. *Trends Biochem. Sci.* 26, 465–468.
- Shoemaker, C.A., Klesius, P.H., 1997. Protective immunity against enteric septicemia in channel catfish, *Ictalurus punctatus* (Rafinesque), following controlled exposure to *Edwardsiella ictaluri*. *J. Fish Dis.* 20, 361–368.
- Shoemaker, C.A., Klesius, P.H., Bricker, J.M., 1999. Efficacy of a modified live *Edwardsiella ictaluri* vaccine in channel catfish as young as seven days post hatch. *Aquaculture* 176, 189–193.
- Shoemaker, C.A., Klesius, P.H., Evans, J.J., 2002. *In ovo* method for utilizing the modified live *Edwardsiella ictaluri* vaccine against enteric septicemia in channel catfish. *Aquaculture* 203, 221–227.
- Shoemaker, C.A., Klesius, P.H., Evans, J.J., 2007. Immunization of eyed channel catfish, *Ictalurus punctatus*, eggs with monovalent *Flavobac-*

- terium columnare vaccine and bivalent *F. columnare* and *Edwardsiella ictaluri* vaccine. Vaccine 25, 1126–1131.
- Shoemaker, R.C., Giclas, P.C., Crowder, C., House, D., Glovsky, M.M., 2008. Complement split products C3a and C4a are early markers of acute Lyme disease in tick bite patients in the United States. Int. Arch. Allergy Immunol. 146, 255–261.
- Singh, A., Singh, I.K., Verma, P.K., 2008. Differential transcript accumulation in *Cicer arietinum* L. in response to a chewing insect *Helicoverpa armigera* and defense regulators correlate with reduced insect performance. J. Exp. Bot. 59, 2379–2392.
- Skirpstunas, R.T., Baldwin, T.J., 2002. *Edwardsiella ictaluri* invasion of IEC-6, Henle 407, fathead minnow and channel catfish enteric epithelial cells. Dis. Aquat. Organ. 51, 161–167.
- Steele-Mortimer, O., St-Louis, M., Olivier, M., Finlay, B.B., 2000. Vacuole acidification is not required for survival of *Salmonella enteric* serovar Typhimurium within cultured macrophages and epithelial cells. Infect. Immun. 68, 5401–5404.
- Sternberg, M.B., Gepstein, S., 2007. Subtractive hybridization techniques to study cellular senescence. Methods Mol. Biol. 371, 289–305.
- Theilacker, C., Sanchez-Carballo, P., Toma, I., Fabretti, F., Sava, I., Kropec, A., Holst, O., Huebner, J., 2009. Glycolipids are involved in biofilm accumulation and prolonged bacteraemia in *Enterococcus faecalis*. Mol. Microbiol. 71, 1055–1069.
- Thune, R.L., Collins, L.A., Pena, M.P., 1997. A comparison of immersion, immersion–oral combination, and injection methods for the vaccination of channel catfish *Ictalurus punctatus* against *Edwardsiella ictaluri*. J. World Aquacult. Soc. 28, 193–201.
- Thune, R.L., Fernandez, D.H., Benoit, J.L., Kelly-Smith, M., Rogge, M.L., Booth, N.J., Landry, C.A., Bologna, R.A., 2007. Signature-tagged mutagenesis of *Edwardsiella ictaluri* identifies virulence-related genes, including a *Salmonella* pathogenicity island 2 class of type III secretion systems. Appl. Environ. Microbiol. 73, 7934–7946.
- Tsujita, T., Tsukada, H., Nakao, M., Oshiumi, H., Matsumoto, M., Seta, T., 2004. Sensing bacterial flagellin by membrane and soluble orthologs of Toll-like receptor 5 in rainbow trout (*Onchorhynchus mikiss*). J. Biol. Chem. 279, 48588–48597.
- Wagner, B.A., Wise, D.J., Khoo, L.H., Terhune, J.S., 2002. The epidemiology of bacterial diseases in food-size channel catfish. J. Aquat. Anim. Health 14, 263–272.
- Wise, D.J., Klesius, P.H., Shoemaker, C.A., Wolters, W.R., 2000. Vaccination of mixed and full-sib families of channel catfish *Ictalurus punctatus* against enteric septicemia of catfish with a live attenuated *Edwardsiella ictaluri* isolate (RE-33). J. World Aquacult. Soc. 31, 206–212.
- Wolters, W.R., Johnson, M.R., 1994. Enteric septicemia resistance in blue catfish and three channel catfish strains. J. Aquat. Anim. Health 6, 329–334.
- Wong, M.L., Medrano, J.F., 2005. Real-time PCR for mRNA quantitation. Biotechniques 39, 75–85.
- Yamagoe, S., Yamakawa, Y., Matsuo, Y., Minowada, J., Mizuno, S., Suzuki, K., 1996. Purification and primary amino acid sequence of a novel neutrophil chemotactic factor LECT2. Immunol. Lett. 52, 9–13.
- Yeh, H.Y., Klesius, P.H., 2007. Molecular cloning and expression of channel catfish, *Ictalurus punctatus*, complement membrane attack complex inhibitor (CD59). Vet. Immun. Immunopath. 120, 246–253.
- Yeh, H.Y., Klesius, P.H., 2008. Complete structure, genomic organization and expression of channel catfish (*Ictalurus punctatus*, Rafinesque 1818) Matrix Metalloproteinase-9 gene. Bios. Biotechnol. Biochem. 72, 702–714.
- Zhang, Y., Arias, C.R., 2007. Identification and characterization of an intervening sequence within the 23S ribosomal RNA genes of *Edwardsiella ictaluri*. Syst. Appl. Microbiol. 30, 93–101.
- Zhang, Y.B., Jiang, J., Chen, Y.D., Zhu, R., Shi, Y., Zhang, Q.Y., Gui, J.F., 2007a. The innate immune response to grass carp hemorrhagic virus (GCHV) in cultured *Carassius auratus* blastulae (CAB) cells. Dev. Comp. Immunol. 31, 232–243.
- Zhang, Y.B., Wang, Y.L., Gui, J.F., 2007b. Identification and characterization of two homologues of interferon-stimulated gene ISG15 in crucian carp. Fish Shellfish Immunol. 23, 52–61.
- Zhao, C.J., Wang, A.R., Shi, Y.J., Wang, L.Q., Liu, W.D., Wang, Z.H., Lu, G.D., 2008. Identification of defense-related genes in rice responding to challenge by *Rhizoctonia solani*. Theor. Appl. Genet. 116, 501–516.
- Zhou, C., Lakso, A.N., Robinson, T.L., Gan, S., 2008. Isolation and characterization of genes associated with shade-induced apple abscission. Mol. Genet. Genomics 280, 83–92.
- Zurovec, M., Dolezal, T., Gazi, M., Pavlova, E., Bryant, P.J., 2002. Adenosine deaminase-related growth factors stimulate cell proliferation in *Drosophila* by depleting extracellular adenosine. Proc. Natl. Acad. Sci. U.S.A. 99, 4403–4408.